

Associations between Seasonal Variation of Heart Rate Variability and Healthy Life Expectancy in Japan

Emi Yuda¹, Masaya Kiso², Yutaka Yoshida³, Norihiro Ueda², and Junichiro Haya²

¹Tohoku University Graduate School of Engineering, Sendai, Japan

²Nagoya City University Graduate School of Medical Sciences, Nagoya, Japan

³Nagoya City University Graduate School of Design and Architecture, Nagoya, Japan

Email: yuda@ieee.org, k.miledge@gmail.com, yoshida@sda.nagoya.ac.jp {nueda,hayano}@med.nagoya.ac.jp

Abstract—Although both summer heat and winter cold increase the risk of health problems, the effects could be modified by living environment and clothes. We investigated the relationship between regional differences in healthy life span (HALE) and seasonal variations in Heart Rate Variability (HRV) that may reflect biological burden caused by heat and cold weather. Using the Allostatic State Mapping by Ambulatory ECG Repository (ALLSTAR) database, we analyzed the 24h standard deviation of sinus rhythm R-R intervals (SDNN) among 86,712 men and 108,771 women across Japan as a measure of HRV. Then, SDNN data were divided into 47 prefectures, averaged over 4 seasons, and associated with the prefectural averages of HALE reported by Japanese government. As the results, in both sexes, prefectural age-adjusted HALE negatively correlated with seasonal variation of prefectural age-adjusted SDNN, while it did not correlate significantly with prefectural age-adjusted SDNN of any seasons. Additionally, there was no significant correlation between prefectural age-adjusted average life expectancy and the seasonal variation of prefectural age-adjusted SDNN. Our observations support the hypothesis that the magnitude of the seasonal variation in biological burden might be a shortening factor for HALE.

Index Terms—ALLSTAR, healthy life expectancy, heart rate variability, seasonal

big data study, we have previously reported that regional differences in HALE but not ALE are associated with regional differences in Heart Rate Variability (HRV); prefectures with higher age-adjusted HRV have longer age-adjusted HALE in both men and women [2]. Using big data of actigraphic physical activity, we have also observed the association between regional difference in adjusted physical activity and age-adjusted HALE [3].

However, it is still unclear why there are differences in HRV and physical activity between prefectures. Although the influence of climate differences between prefectures can be considered, the regional distribution of HALE, HRV, and physical activity observed so far does not match the distribution of average temperature and precipitation. In this study, we hypothesized that the physiological effects of seasonal variations may be a factor of regional variation in HALE. Although both summer heat and winter cold increase the risk of health problems, the effects could be modified by living environment and clothes. We therefore investigated the seasonal variations in HRV that may reflect biological burden caused by heat and cold weather. Using an ambulatory ECG big data, Allostatic State Mapping by Ambulatory ECG Repository (ALLSTAR) [4], we analyzed the relationship between regional differences in HALE and seasonal variations in HRV.

I. INTRODUCTION

Along with increasing aging societies, the gap between Average Life Expectancy (ALE) and healthy life expectancy (HALE) became a critical issue (Fig. 1). For Japanese men, the last 9.79 years of their lives cannot live on their own for health reasons. This period is 12.99 year in Japanese women. By improving preventive medicine and improving lifestyle through public health awareness, HALE has gradually increased, but the gap has hardly narrowed as ALE has also increased.

It is recognized that HALE shows a distinct regional difference in Japan [1]. Although the clear reasons for the regional difference are known, in an ambulatory ECG

II. METHODS

A. Database

We studied Holter ECG big data that has been accumulated by ALLSTAR Research Group since 2009. The database consists of >300 thousand 24h Holter ECG recorded with standardized 1-lead recorders (Cardy 303 pic α , Suzuken CO., LTD, Nagoya, Japan) between April 2012 and July 2014 in almost all prefectures in Japan. We used data only from subjects >20 whose electrocardiogram showed sinus rhythm >90% of the recording length with no abnormal finding on clinical ECG diagnosis. The study protocol has been approved by the Research Ethics Committee of Nagoya City University Graduate School of Medical Sciences (No. 709).

Manuscript received February 24, 2020; revised May 29, 2020.





Figure 1. Average life expectancy (ALE) and healthy life expectancy (HALE) in Japanese men and women (from Japanese Health, Labor, and Welfare Ministry).

B. Data Analysis

The Holter ECG signals were analyzed by a Holter ECG scanner (Cardy 5, Suzuken Co., LTD, Nagoya, Japan), on which all QRS waves were detected and labeled for rhythm annotations. We analyzed 24-hour SD only from normal-to-normal RR intervals (SDNN) as an HRV measure among 86,712 men (median age [IQR], 68 [56-71] yr) and 108,771 women (70 [58-79] yr).

SDNN were divided into the matrix of 47 prefectures, four seasons, and least square means adjusted for the effect of age were calculated for each cell. The seasonal difference in SDNN of each prefecture was defined as the range of difference among four seasons (max) and was associated with the prefectural averages of HALE and ALE reported by Japanese government (the prefecture ranking of HALE published by the Japanese Health, Labour, Welfare Ministry [1] (Fig. 2). Only prefectures with an enough sample size (150 for each sex) were used for analysis.





Figure 2. Seasonal difference in SDNN and healthy life expectancy (HALE) in Japanese prefectures. SDNN dif = range of seasonal difference in SDNN; NSD = no sufficient data.

Figure 3. Annual average temperature and seasonal difference in SDNN in Japanese prefectures. SDNN dif = range of seasonal difference in SDNN; NSD = no sufficient data.

III. RESULTS

Although there were no significant correlations between HALE and SDNN of any seasons, significant negative correlation between seasonal difference in SDNN and HALE was observed in men, but not in women (Table I). In men, there was a tendency that seasonal difference in SDNN was larger in prefectures where HALE was shorter than the median (seasonal SDNN difference, 10.1 ± 5.7 vs. 7.3 ± 3.8 ms, $P = 0.08$; Fig. 2). In neither sex, any seasonal SDNN nor seasonal change in SDNN correlated with ALE (Fig. 3).

IV. DISCUSSIONS

In this study, we analyzed the relationship between seasonal variation in SDNN and HALE among Japanese prefectures. We observed the negative association between seasonal difference in SDNN and HALE only in men but not in women.

Although the presence of seasonal differences in HRV has been reported [6], the interaction between the effects of seasons and occupation has also been recognized. The contrast of HRV between summer and winter may be greater in outdoor workers than indoor workers. This suggests that the health burden of seasonal environmental changes may be related not only to seasonal variations in the outdoor climate environment but also to seasonal variations in the indoor living environment. On the other hand, buildings and clothes in cold regions have high cold protection functions, but those functions are low in warm regions. Conversely, cold regions have a weak function against heat than warm regions. Consequently, health burden in winter season could be greater in warm regions and that in summer season could be greater in cold regions. These suggest that seasonal health effects may depend on many factors including regional climate, occupation, home and office architecture, and clothes customs.

In the present study, we observed significant association between seasonal difference in SDNN and HALE only in men but not in women (Fig. 2). This may not be attributed to physiological gender difference, because we previously observed significant association between the prefectural average of age-adjusted SDNN and prefectural average of HALE in both men and women [2]. In general, men still have more opportunities to spend outside than women in Japan. Thus, current observations suggest the potential impact of differences in the causes of seasonal variations in SDNN, i.e., whether due to seasonal variations in the outdoor climate or seasonal variations in the indoor environment. We also observed the significant association between annual average temperature and seasonal variation in SDNN in either sex (Fig. 3). This observation supports that seasonal health effects are explained only by outdoor climate. Fig. 3 also shows that there are considerable differences in the seasonal variation of SDNN even within cold, medium and warm regions, suggesting the impact of unmeasured factors such as economy and industry. These points need further analyses in future studies.

This study has limitations. First, the seasonal variation in HRV was estimated by comparing the data obtained in different seasons recorded in different subjects. Although many subjects in each season can be expected to average out the effects of individual differences, it is ideal to compare data obtained in different seasons in the same subject. Second, the ALLSTAR database consisted of Holter ECG recorded for clinical reasons. Although we used only data with no abnormal finding on clinical ECG diagnosis, the influence cannot be excluded for diseases that do not show abnormalities in ECG particularly, those with seasonal variations in incidence. Third, we compared prefectural average of HALE and prefectural average of HRV. While HALE reflects the value expected of people aged 0 HRV was measured in adult >20 years and most of them were in their 50s to 70s. Although we adjusted the effect of age on HRV, the actual healthy life span of the subjects could differ from the HALE used in this study.

V. CONCLUSIONS

Using ALLSTAR big data, we analyzed the relationship between seasonal variation in SDNN and HALE among Japanese prefectures. We observed that HALE in men was shorter in prefectures with greater seasonal variation in HRV, although no significant association was observed for women. The magnitude of the seasonal variation in HRV may be a shortening factor for HALE in men but not women.

CONFLICT OF INTEREST

The authors declare no conflict of interest

AUTHOR CONTRIBUTIONS

EY managed the database, conceived the study, and critically reviewed the manuscript; MK maintained the database and analyzed data; YY analyzed the data; NU managed the database and critically reviewed the manuscript; JH analyzed the data and wrote the manuscript; all authors had approved the final version.

ACKNOWLEDGMENT

The authors would like to thank the ALLSTAR project committee and Suzuken Co., LTD. for their support for this study.

REFERENCES

- [1] Progress of each goal in Healthy Japan 21 (2nd stage). (2019) The Ministry of Health, Welfare, and Labour. [Online]. Available: <http://www.mhlw.go.jp/file/05-Shingikai-10601000-Daijinkanboukouseikagakuka-Kouseikagakuka/sinntyoku.pdf>
- [2] E. Yuda, Y. Furukawa, Y. Yoshida, J. Hayano, and ALLSTAR Investigators, "Association between regional difference in heart rate variability and inter-prefecture ranking of healthy life expectancy: ALLSTAR Big Data Project in Japan," in *Big Data Technologies and Applications: Proceedings of the 7th EAI International Conference, BDTA 2016*, J. J. Jung and P. Kim Eds. Seoul, Korea: Springer Nature, 2017, pp. 23-28.
- [3] E. Yuda, Y. Yoshida, H. Ogasawara, J. Hayano, and ALLSTAR Study Group, "Ambulatory physical activity monitoring to know

healthy life expectancy,” in *Proc. 8th International Workshop on Biosignal Interpretation (BSI2016)*, Y. Yamamoto, T. Nomura, S. Cerutti, H. Dickhaus, and K. Yana, Eds., Osaka, Japan, Nov. 2016, pp. 1-3.

- [4] J. Hayano, K. Kiyono, E. Yuda, Y. Yamamoto, and I. Kodama. (2018). Holter ECG big data project: Allostatic state mapping by ambulatory ECG repository (ALLSTAR). *International Journal of Information Research and Review*. [Online]. 5(7). pp. 5617-5624. Available: <https://www.ijirr.com/holter-ecg-big-data-project-allostatic-state-mapping-ambulatory-ecg-repository-allstar>.
- [5] ALLSTAR Research Group. (2019). Allostatic state mapping by ambulatory ECG repository (ALLSTAR). [Online]. Available: <http://www.med.nagoya-cu.ac.jp/mededu.dir/allstar/index.html>
- [6] J. Kristiansen, A. Olsen, J. H. Skotte, and A. H. Garde, “Reproducibility and seasonal variation of ambulatory short-term heart rate variability in healthy subjects during a self-selected rest period and during sleep,” *Scand. J. Clin. Lab. Invest.*, vol. 69, no. 6, pp. 651-61, 2009.
- [7] A. Markov, I. Solonin, and E. Bojko, “Heart rate variability in workers of various professions in contrasting seasons of the year,” *Int. J. Occup. Med. Environ. Health*, vol. 29, no. 5, pp. 793-800, 2016.

Copyright © 2020 by the authors. This is an open access article distributed under the Creative Commons Attribution License ([CC BY-NC-ND 4.0](https://creativecommons.org/licenses/by-nc-nd/4.0/)), which permits use, distribution and reproduction in any medium, provided that the article is properly cited, the use is non-commercial and no modifications or adaptations are made.



Emi Yuda was born in Tokyo, Japan. She studied informatics at M. V. Lomonosov Moscow State University until 2003 and then received M.S. degree from Tsukuba University, Japan. She received Ph.D. from Nihon University in 2019. From 2013 to 2014 she was a research assistant at Santa Monica College in California, USA. From 2015 to 2019, she was a NEDO project researcher in Nagoya City University Graduate School of Medical Sciences. Since 2019, she has been an

assistant professor in Tohoku University Graduate School of Engineering. Her currently research is Medical Informatics, Data Science and Bio-signal Processing. She has many achievements in field of Informatics and Big Data.



Masaya Kisoehara graduated Nagoya City University Medical School, Nagoya, Japan and received M.D. degree in 2014. From 2012 to 2013, he received research training for biosignal processing (analysis of heart rate variability) at Nagoya City University Medical School. In 2016, he has received residency training of radiological diagnosis and interventional radiology in Nagoya City University Hospital. Since 2018, he is a student of Medicine at Nagoya City University

Graduate School of Medical Sciences. His current interests are application to medical treatment include image diagnosis with artificial intelligence.



Yutaka Yoshida studied the business administration and computer science at Aichi Institute of Technology and received Ph.D. degree in 2008. He was a project researcher at Knowledge Hub of Aichi from 2011 to 2015 and was a researcher at Nagoya City University Graduate School of Medical Sciences from 2016 to 2017. Since 2018, he has been a researcher at Nagoya City University Graduate School of Design and Architecture. His

specialized field is biological information engineering, signal processing and ergonomics. He received the paper award at the Japan Society of Neurovegetative Research in 2005 and 2007.



Junichiro Hayano graduated Nagoya City University Medical School, Nagoya, Japan and received M.D. degree in 1980. From 1981 to 1983, he received residency trainings of psychosomatic medicine in Kyushu University School of Medicine, Fukuoka, Japan. He obtained Ph.D. degree (Dr. of Medical Science) in 1988 from Nagoya City University Graduate School of Medical Sciences. From 1990 to 1991, he was a visiting associate at the Behavioral Medicine Research Center, Duke

University Medical Center, Durham, NC, USA. In 1984, he got a faculty position at Nagoya City University Medical School and has been a Professor of Medicine at Nagoya City University Graduate School of Medical Sciences since 2003. His current interests are applications of bio-signal processing to health sciences.