

Enhancement of Algal Biomass Accumulation Using Undiluted Anaerobic Digestate

Juliana G. Ivanova¹, Ivanina A. Vasileva², Lyudmila V. Kabaivanova³

¹Department of Experimental Algology, Institute of Plant Physiology and Genetics, Bulgarian Academy of Sciences, Sofia, Bulgaria

²Institute of Plant Physiology and Genetics, Bulgarian Academy of Sciences, Sofia, Bulgaria

³Department of Applied Microbiology, The Stephan Angeloff Institute of Microbiology, Bulgarian Academy of Sciences, Sofia, Bulgaria

Email: ivanina_vasileva1@abv.bg, {juivanova, lkabaivanova}@yahoo.com

Abstract—Microalgae are unicellular photosynthetic organisms with industrial and economic perspectives. Their cultivation has a great deal of potential for many applications. Microalgae possess high growth and nutrient uptake rates, carbon capture and biomass generation. The cultivation of microalgae needs to be improved in order to reduce the cost of the produced biomass. Anaerobic digestion is a waste-to-energy technology employing a consortium of anaerobic microorganisms and resulting in continuous production of methane-rich biogas, with intermittent release of effluent digestate rich in undigested solids, organic and inorganic compounds and metal salts. The results in this work demonstrate the possibility of improving algal biomass accumulation using different anaerobic digestates and suitable algal strains. We investigated the growth and development of seven microalgal species – green, blue-green, red and yellow-green in the resultant digestate from methane production. The digestate was decolorized with active charcoal and used as medium in non-diluted and diluted with water (1:1) form. Most abundant growth was registered after two weeks of extensive cultivation for the green algae *Scenedesmus acutus*, followed by *Klebsormidium flaccidum*, and yellow-green *Trachydiscus minutus*. Further intensive cultivation of *Scenedesmus acutus* was performed and good growth was confirmed by estimating the average specific growth rate ($\mu=0.30\text{ d}^{-1}$) and pigment accumulation (total pigment content 7.45% of DW). Our results showed that the digestate, obtained straight after a biomethane generating anaerobic process, with no additional water, is a suitable cultivation medium. Compared to the control with a standard medium, it enhanced the growth of *Scenedesmus acutus* 6.5 times in one week. This is a good precondition for a reliable application of this digestate in practice for production of algal biomass with its metabolites, instead of being released as a waste.

Index Terms—microalgae, digestate, growth rate, pigments, *Scenedesmus acutus*

I. INTRODUCTION

Microalgae are fast-growing phototrophic microorganisms which utilize light energy and inorganic

nutrients (carbon dioxide, nitrogen, phosphorus etc.) [1] And in turn synthesize valuable biomass compounds, such as lipids, proteins, carbohydrate and pigments [2], [3]. These compounds find application in various industries: pharmaceutical, food industry, agriculture, green chemistry and the bioenergy sectors [4]. This makes the reduction of the price of the biomass production essential for the further development of the algal biotechnology. Among the major costs for algal cultivation are the ones for nutrients, such as nitrogen and phosphorous. These components could be found in large quantities in wastewaters from different industries [5], [6]. Microalgae offer an elegant solution to tertiary and quaternary treatments due to their ability to use inorganic nitrogen and phosphorus for their growth [7], which is supported by their high capacity of inorganic nutrient uptake [8].

In the last decades, biogas production has developed rapidly. The major fraction in this anaerobic fermentation is the liquid digestate. Typically, anaerobic digestate is characterized by high levels of macro and microelements [9]. As the amount of nutrients in the digestate cannot be significantly reduced during the anaerobic digestion [10], it needs to be effectively recycled afterwards. This leads to the idea of utilizing the product by microalgae instead of discharging it into the environment [11], [12]. There are reports of successful cultivation of microalgae such as *Chlorella*, *Scenedesmus*, *Phaeodactylum* and *Pavlova* in anaerobic digestate [13]–[15].

Selection of appropriate strains is critical for producing valuable biomass [16]. Examples of targeted biosynthesis are the species *Klebsormidium* and *Stigeoclonium*, which accumulate valuable unsaturated fatty acids such as C18: 2 and C18: 3 in large quantities [17]. A main requirement for sustainable digestate treatment is a high biomass accumulation, therefore, the next step is the selection of suitable conditions for algae cultivation [6]. It is well known that each microalgal strain has its favorable growth conditions, such as pH, light, temperature, salinity, and the preferred nitrogen form and N / P ratio [2], [18]–[20]. The aim of this work was to find a perspective algal strain that grows well on

undiluted digestate from anaerobic digestion of agro-waste during biogas production. This study presents a possible solution for reducing the costs of algal cultivation, and production of biomass and valuable metabolites.

II. MATERIALS AND METHODS

A. Biological Material and Cultivation Conditions

For our experiments, seven microalgal strains: green algae *Scenedesmus acutus*, *Scenedesmus* sp. BGP, *Klebsormidium flaccidum* and *Coelastrrella* sp. BGV, yellow-green alga *Trachydiscus minutus*, red alga *Porphyridium aerugineum*, blue-green alga *Synechocystis salina* were used. All of the abovementioned algae are deposited in the collection of Laboratory of Experimental Algology, Institute of Plant Physiology and Genetics, BAS, Sofia, Bulgaria.

B. Extensive Cultivation

Extensive algae cultivation was carried out at room temperature and natural daylight for a period of 14 days. A standard culture medium of Setlik, modified by [21], was used as a control. Three variants of anaerobic digestates from a methanogenic laboratory bioreactors were tested as nutrient media for algal cultivation: Variant 1: digestate from a mesophilic anaerobic process (37 °C) with substrate wheat straw 10 g L⁻¹, chemically pretreated; Variant 2: digestate with substrate wheat straw (37 °C) and Variant 3: digestate from a thermophilic process (55 °C) with substrate wheat straw. The liquid digestate was subjected to clarification by adsorption with 8g 100mL⁻¹ active charcoal (Flika) overnight at room temperature for all the three variants.

1) Chemical pretreatment

The digestate was treated with 28% NH₄OH, polyethylene glycol-4000 (3%) and water in a ratio of 1:0.5:20: NH₄OH: H₂O. The mixture underwent heating at 90 °C for 5h in a water bath. This was followed by rinsing till neutral pH.

a) Cell count

To evaluate the growth and development of the investigated strain, cell count was carried out using a Burk counting chamber.

b) Elemental analysis

Elemental composition of the obtained digestate was performed with EuroEA 3000 automatic analyzer.

C. Intensive Cultivation

An initial algal culture density of 0.8 mg. mL⁻¹ dry weight (DW) was used for all experiments. Cultivation was carried out at 25 °C and continuous illumination (132 μmol photons m⁻² s⁻¹). A carbon source was provided by bubbling sterile 2% CO₂ (v/v). The standard culture medium of Setlik [21], was used for the control cultivation. Variant 1 of the digestate was used in non-diluted and 50% diluted form for all the treatments.

1) Dry weight and specific growth rate

The growth was evaluated by the increase in algal biomass. The DW (mg mL⁻¹) was determined gravimetrically. Algal suspensions (3×5 mL each) were

filtered through Whatman GF/C glass filters (Whatman International Ltd, Maidstone, UK), rinsed with tap water to eliminate salts and oven dried at 105 °C till a constant weight. The specific growth rate [μ] was calculated using the following formula:

$$\mu = \ln(m_2/m_1) / (t_2 - t_1) \quad [22],$$

where m_t are the dry weights at the different days ($t_1 = 0$ and $t_2 = 6$).

2) Pigment content

Pigments - chlorophyll a, chlorophyll b, carotenoids, were measured spectrophotometrically at 665 nm, 645 nm and 460 nm, respectively, using a T70 UV/Vis (PG Instruments Ltd, Leicester, UK) spectrophotometer after extraction with boiling methanol. Using the absorptions, the pigment content was calculated, employing the Mackiney formulas [23].

D. Data Analysis A

The experiments were conducted in triplicate. The data were presented as the means \pm SD. The difference between the treatments was statistically analyzed by oneway ANOVA followed by the Bonferroni's post hoc test at significance level of $p < 0.05$, using GraphPAD InStat Software (San Diego, CA, USA).

III. RESULTS AND DISCUSSION

A. Extensive Cultivation of Microalgae on Digestate

In order to investigate the growth abilities of microalgal strains in anaerobic digestate as a medium, batch experiments were conducted at extensive cultivation conditions for a period of 14 days. On Fig.1 is presented the growth of the 7 strains in Variant 1 of digestate, determined by the increased number of algal cells for all the strains. The graph shows that most of the algae were growing well on this waste product. The representatives of green microalgae showed the highest increase. The maximal growth was registered for *Scenedesmus acutus*, followed by *Klebsormidium flaccidum* ($p > 0.05$). Good growth was also observed for the eustigmatophycean alga *Trachydiscus minutus* ($p < 0.05$). Blue-green and red microalgae showed the slowest growth.




Figure 1. Growth of 7 different microalgal species on digestate from a mesophilic metanogenic process. Means with different lowercase letters are significantly different ($P < 0.05$) between the different algal strains.

Therefore, the subsequent experiments were performed with the alga that showed the greatest increase in cell mass (about 25-fold) - *Scenedesmus acutus*. It was grown in all the three variants of the anaerobic digestate, described in the Materials and methods section.

The results showed that Variant 1 of the digestate is the most suitable cultivation medium (Fig. 2). According to the cell count, there was no significant difference in the growth of *Scenedesmus acutus* in this medium compared to the control variant ($p>0.05$) at the end of the cultivation. Slightly lower algal growth was observed in Variant 2 followed by Variant 3. The elemental analysis indicated that when Variant 1 was used, the assimilation of N, H and C was also the highest (Table I).




Figure 2. Growth of *Scenedesmus acutus* in different digestate variant.

TABLE I. ELEMENTAL ANALYSIS OF DIGESTATES

| | Carbon, % | Nitrogen, % | Hydrogen, % |
|-----------|------------|-------------|-------------|
| Control | 12.70±0.97 | 2.79±0.16 | 2.33±0.21 |
| Variant 1 | 7.68±0.54 | 1.65±0.11 | 1.49±0.11 |
| Variant 2 | 12.33±1.04 | 2.70±0.23 | 2.41±0.19 |
| Variant 3 | 8.57±0.63 | 1.86±0.17 | 1.86±0.15 |

B. Intensive Cultivation

Based on this, we used Variant 1 of the digestate (in non-diluted and 50% diluted form) as a cultivation medium for the intensive cultivation of the chosen alga – *Scenedesmus acutus* (Fig. 3, Fig. 4).

The results acquired were promising for both diluted and non-diluted digestate. Over the 144-hour cultivation period, the DW of the cultures grown in the undiluted digestate reached values of $5.2 \pm 0.2 \text{ mg mL}^{-1}$, even exceeding the amount of accumulated biomass in the standard medium (control sample).



Figure 3. Intensive cultivation: at the beginning (left photo) and after 6 days of cultivation (right photo) in 100% digestate - first tube and 50% digestate - second tube.




Figure 4. Intensive growth of *Scenedesmus acutus* in Variant 1 of digestate (100% and 50%). * $p < 0.05$ indicates significant difference from control values.

Following the dynamics of the process, initially (till 72h) a higher increase in the variant with 50% diluted digestate was registered (Fig. 4). In our opinion, this is due to the greater clarity of the diluted digestate at the beginning of the cultivation period. At the end of the exponential phase the elevated optical density due to the increased number of cells and the depleted nutrients, lead to decreased growth rate of *Scenedesmus* in this sample (Fig. 4).

Fig. 5 illustrates the average growth rate over the whole period. The highest average $\mu \text{ (d}^{-1}\text{)}$ was recorded in the sample with undiluted digestate (0.30 d^{-1}) followed by the control (0.29 d^{-1}) and the variant with 50% dilution (0.28 d^{-1}).




Figure 5. Specific growth rate of *Scenedesmus acutus* in digestate.

C. Pigment Analyses

The good growth and development of the microalgal culture of *Scenedesmus acutus* was proved by estimation of its pigment content.

The absorbance spectra peaks were obtained from methanolic extracts of intensively cultivated algal samples (Fig. 6). The results for the pigment composition of *Scenedesmus acutus* (Fig. 6, Table I) correlated with the growth of the algae and the increasing biomass production, which were highest in the samples, cultivated in the undiluted digestate.

Table II illustrates the results for the quantitative composition of the pigments from the samples, harvested at the 144th h of the intensive cultivation of *Scenedesmus acutus*. The values for chlorophylls and carotenoids are similar for control and undiluted variants (Table II).




Figure 6. Spectrophotometric estimation of absorption of pigments at the end of intensive cultivation. Green line-sample 2, grown in 50% diluted digestate; violet –control sample, grown in standard medium; pink – sample 1, grown in undiluted digestate.

TABLE II PIGMENT CONTENT (% OF DW)

| | chl. "a" | chl. "b" | carotenoids |
|----------------|-----------|-----------|-------------|
| control | 2.8 ± 0.1 | 1.2 ± 0.2 | 3.15 ± 0.37 |
| 100% digestate | 2.9 ± 0.1 | 1.3 ± 0.2 | 3.25 ± 0.28 |
| 50% digestate | 2.1 ± 0.1 | 0.8 ± 0.1 | 2.5 ± 0.16 |

The results obtained for the best growth in digestate for the green algae are confirmed by other authors [24], [25].

So far, most studies for growth of algae in digestate have been performed with large dilutions of 5 to 30 times [12], [24]. This reduces the efficiency many times, including addition of pure water in the cultivation process. In our work, we used activated charcoal to reduce the color of the digestate, thus were able to achieve substantial growth both with 50% dilution and without dilution (Fig. 4 and Fig. 5). At the end of the exponential phase in the non-diluted digestate was registered better growth of *Scenedesmus acutus*, about 1.16 times higher than diluted variant.

In the study of [26] was reported that the average specific growth rates of *Chlorella* and *Micractinium* are about 0.13 d⁻¹ and 0.14 d⁻¹. The scientific team of [12] obtained a specific growth rate of 0.24 d⁻¹ for the alga *Phaeodactylum tricornutum*. Our results showed μ of 0.30 (d⁻¹) on average over the entire cultivation period for the undiluted variant, which is 1.25 times higher than the reported by [26]. It should be noted that neither during the extensive, nor the intensive cultivation, any additional components were added to the culture medium.

Reference [27] reported an achieved yield of algal biomass in the range of 1.7 to 2.1 mg mL⁻¹ after 8 days of intensive cultivation of *Chlorella* sp. in 50% diluted anaerobically digested sludge. Our results are two times higher, as over a 6-day cultivation period the DW of *Scenedesmus acutus* was 5.2±0.2 mg mL⁻¹ without dilution, and 4.5±0.3 mg mL⁻¹ with 50% dilution. This result is in agreement with a previous study of [28], claiming that the strains associated with this species were

most effective when grown with substrates derived from waste water.

IV. CONCLUSION

The approach combining algal biomass production with utilization of digestate from a methane generating process creates a cost-effective management of an integrated biotechnological process. Moreover, the use of microalgae for waste remediation has an additional advantage. We proved the possibility of enhancing algal biomass production choosing the appropriate conditions for the anaerobic digestion process, as well as for the extensive and intensive cultivation of the algal strains.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

AUTHOR CONTRIBUTIONS

J. I. and L. K. planned the project; L. K. provided the digestate for the experiments; J. I. and I. V. conducted the experiments, analyzed the data and wrote the article. All authors had approved the final version.

ACKNOWLEDGMENT

This work was supported by a grant from the Bulgarian Science Fund, project KII-06-H26/5.

REFERENCES

- [1] N. Kobayashi, E. A. Noell, A. Barnesc, A. Watsonc, J. N. Rosenbergd, G. Ericksonc, and G. A. Oylerad, "Characterization

- of three *Chlorella sorokianiana* strains in anaerobic digested effluent from cattle manure," *Bioresour. Technol.*, vol. 150, pp. 377-386, December 2013.
- [2] G. Markou and D. Georgakakis, "Cultivation of filamentous cyanobacteria (bluegreen algae) in agro-industrial wastes and wastewaters: A review," *Applied Energy*, vol. 88, pp. 3389-3401, October 2011.
 - [3] R. S. Gours, A. Kant, and R. S. Chauhan, "Screening of micro algae for growth and lipid accumulation properties," *J. Algal Biomass Utiln.*, vol. 5, pp. 38-46, 2014.
 - [4] O. Pulz and W. Gross, "Valuable products from biotechnology of microalgae," *Appl. Microbiol. Biotechnol.*, vol. 65, pp. 635-648, November 2004.
 - [5] J. Liu, B. Danneels, P. Vanormelingen, and W. Vyverman, "Nutrient removal from horticultural wastewater by benthic filamentous algae *Klebsormidium* sp., *Stigeoclonium* spp. and their communities: From laboratory flask to outdoor Algal Turf Scrubber (ATS)," *Water Res.*, vol. 92, pp. 61-68, April 2016.
 - [6] S. Van den Hende, V. Beelen, G. Bore, N. Boon, and H. Vervaeren, "Up-scaling aquaculture wastewater treatment by microalgal bacterial flocs: from lab reactors to an outdoor raceway pond," *Bioresour. Technol.*, vol. 159, pp. 342-354, May 2014.
 - [7] W. Mulbry, P. Kangas, and S. Kondrad, "Toward scrubbing the bay: nutrient removal using small algal turf scrubbers on Chesapeake Bay tributaries," *Ecol. Eng.*, vol. 36, no. 4, pp. 536-541, April 2010.
 - [8] R. Blier, G. Laliberte, and J. De la Noue, "Tertiary treatment of cheese factory anaerobic effluent with *Phormidium bohneri* and *Micractinium pusillum*," *Bioresour. Technol.*, vol. 52, pp. 151-155, 1995.
 - [9] A. Xia and J. D. Murphy, "Microalgal cultivation in treating, liquid digestate from biogas systems," *Trends Biotechnol.*, vol. 34, no. 4, pp. 264-275, April 2016.
 - [10] M. Franchino, E. Comino, F. Bonaa, and V. A. Riggio, "Growth of three microalgae strains and nutrient removal from an agro-zootechnical digestate," *Chemosphere*, vol. 92, no. 6, pp. 738-744, July 2013.
 - [11] A. Husam, A. Hajar, R. G. Riefler, and B. J. Stuart, "Anaerobic digestate as a nutrient medium for the growth of the green microalga *Neochloris oleoabundans*," *Environ. Eng. Res.*, vol. 21, no. 3, pp. 265-275, September 2016.
 - [12] D. Veronesia, A. Idaa, G. D'Imporzano, and F. Adani, "Chemical microalgae cultivation: nutrient recovery from digestate for producing algae biomass," *Eng. Transact.*, vol. 43, pp. 2283-9216, May 2015.
 - [13] H. J. Choi and S. M. Lee, "Effects of microalgae on the removal of nutrients from wastewater: Various concentrations of *Chlorella vulgaris*," *Environ. Eng. Res.*, vol. 17, pp. S3-S8, December 2012.
 - [14] M. Ras, L. Lardon, S. Bruno, N. Bernet, and J. Steyer, "Experimental study on a coupled process of production and anaerobic digestion of *Chlorella vulgaris*," *Bioresour. Technol.*, vol. 102, pp. 200-206, January 2011.
 - [15] R. Tao, V. Kinnunen, R. Praveenkumar, A. M. Lakaniemi, and J. A. Rintala, "Comparison of *Scenedesmus acuminatus* and *Chlorella vulgaris* cultivation in liquid digestates from anaerobic digestion of pulp and paper industry and municipal wastewater treatment sludge," *J. Appl. Phycol.*, vol. 29, no. 6, pp. 2845-2856, December 2017.
 - [16] G. Roeselers, M. C. M. van Loosdrecht, and G. Muyzer, "Phototrophic biofilms and their potential applications," *J. Appl. Phycol.*, vol. 20, no. 3, pp. 227-235, June 2008.
 - [17] J. Liu, P. Vanormelingen, and W. Vyverman, "Fatty acid profiles of four filamentous green algae under varying culture conditions," *Bioresour. Technol.*, vol. 200, pp. 1080-1084, January 2016.
 - [18] A. Besson and P. Guiraud, "High-pH-induced flocculation-flotation of the hypersaline microalga *Dunaliella salina*," *Bioresour. Technol.*, vol. 147, pp. 464-470, November 2013.
 - [19] J. Liu and W. Vyverman, "Differences in nutrient uptake capacity of the benthic filamentous algae *Cladophora* sp., *Klebsormidium* sp. and *Pseudanabaena* sp. under varying N/P conditions," *Bioresour. Technol.*, vol. 179, pp. 234-242, March 2015.
 - [20] T. Cai, S. Y. Park, and Y. Li, "Nutrient recovery from wastewater streams by microalgae: Status and prospects," *Renew. Sustain. Energy Rev.*, vol. 19, pp. 360-369, March 2013.
 - [21] D. Georgiev, H. Dilov, and S. Avramova, "Millieu nutritif tamponne et méthode de culture intensive des microalgues vertes," *Hydrobiol.*, vol. 7, pp. 14-23, July 1978.
 - [22] M. Levasseur, P. A. Thompson, and P. J. Harrison, "Physiological acclimation of marine phytoplankton to different nitrogen sources," *J. Phycol.*, vol. 29, no. 5, pp. 587-595, October 1993.
 - [23] G. Mackinney, "Criteria for purity of chlorophyll preparations," *J. Biol. Chem.*, vol. 132, pp. 91-109, January 1940.
 - [24] L. Zuliani, N. Frison, A. Jelic, F. Fatone, D. Bolzonella, and M. Ballottari, "Microalgae cultivation on anaerobic digestate of municipal wastewater, sewage sludge and agro-waste," *Int. J. Mol. Sci.*, vol. 17, no. 10, pp. 1692-1696, October 2016.
 - [25] R. Chen, R. Li, L. Deitz, Y. Liu, R. J. Stevenson, and W. Lao, "Freshwater algal cultivation with animal waste for nutrient removal and biomass production," *Biomass. Bioenerg.*, vol. 39, pp. 128-138, October 2012.
 - [26] W. M. Wang, W. C. Kuo-Dahab, S. Dolan, and C. Park, "Kinetics of nutrient removal and expression of extracellular polymeric substances of the microalgae, *Chlorella* sp. and *Micractinium* sp., in wastewater treatment," *Bioresour. Technol.*, vol. 154, pp. 131-137, February 2014.
 - [27] A. M. Åkerström, L. M. Mortensen, B. Rusten, and H. R. Gíslersdóttir, "Biomass production and nutrient removal by *Chlorella* sp. as affected by sludge liquor concentration," *J. Environ. Manage.*, vol. 144, pp. 118-124, November 2014.
 - [28] P. Bohutskyi, K. Liu, L. K. Nasr, N. Byers, J. N. Rosenberg, G. A. Oyler, M. J. Betenbaugh, and E. J. Bouwer, "Bioprospecting of microalgae for integrated biomass production and phytoremediation of unsterilized wastewater and anaerobic digestion centrate," *Appl. Microbiol. Biotechnol.*, vol. 99, no. 14, pp. 6139-6154, July 2015.

Copyright © 2020 by the authors. This is an open access article distributed under the Creative Commons Attribution License ([CC BY-NC-ND 4.0](https://creativecommons.org/licenses/by-nc-nd/4.0/)), which permits use, distribution and reproduction in any medium, provided that the article is properly cited, the use is non-commercial and no modifications or adaptations are made.



Juliana G. Ivanova graduated from Sofia University "St. Kliment Ohridski", Faculty of Biology, with a M.Sc. degree in Biotechnology. Dr. Ivanova received her Ph.D. degree in Plant Physiology at the Institute of Plant Physiology and Genetics, Bulgarian Academy of Sciences. She is currently an Assistant Professor at the same Institute. Professional interests: Algae cultivation, immobilization, interactions

between bacteria and algae, biologically active substances from algae, purification of water and soils from heavy metals and organic pollutants.



Ivanina A. Vasileva received her M.Sc. degree in Plant Biotechnologies at Sofia University "St. Kliment Ohridski", Sofia, Bulgaria in 2015. Her PhD degree in biochemistry was acquired at the Institute of Plant Physiology and Genetics, Bulgarian Academy of Sciences, Sofia, Bulgaria in 2019. She currently works as an Assistant Professor at the Institute of Plant Physiology and Genetics, Bulgarian Academy of Sciences. Dr. Vasileva's professional interests are focused on physiology of algae, algal proteins, biologically active substances derived from algae.



Lyudmila V. Kabaivanova graduated from Sofia University “St. Kliment Ohridski”, Faculty of Biology, with a MSc degree in Biochemistry and Microbiology - 1992. She received her PhD degree in Microbiology at the “Stephan Angeloff” Institute of Microbiology, Bulgarian Academy of Sciences in 2005. She is currently Associate Professor (since 2011), scientific secretary of the same institute and Head of Department “Applied microbiology”. Her main scientific interests are in the field of biodegradation, bioremediation, biologically active substances from bacteria and algae.