

Phytochemical Screening and Evaluation of Antioxidant, Antibacterial Activities of Ethanol Extract from *Combretum quadrangulare* Collected in Vietnam

Tran Thanh Men^{1,*}, Huynh Kim Yen², Huynh Thi Cam La³, Nguyen Hoang Son⁴, Do Tan Khang⁵ and Nguyen Trong Tuan¹

¹Department of Biology, College of Natural Sciences, Can Tho University, Cantho City 90000, Vietnam

²Department of Biology, College of Agriculture and Rural Development, Kien Giang University, Kien Giang Province 91000, Vietnam

³Department of Chemistry, Faculty of Basic Sciences, Vinh Long University of Technology Education, Vinh Long City, 890000 Vietnam

⁴Department of Molecular Biotechnology, Biotechnology Research and Development Institute, Can Tho University, Cantho City, 94000, Vietnam

⁵Department of Chemistry, College of Natural Sciences, Can Tho University, Cantho City 900000, Vietnam
Email: {ttmen, lanb1704486, dtkhang, trongtuan}@ctu.edu.vn, hkyen@vnkgu.edu.vn, nghson92@gmail.com

Abstract *Combretum quadrangulare* Kurz has long been considered a precious herb, which is very useful in treating various kinds of diseases on humans and aquatic animals. The study was conducted to evaluate the antioxidant and antibacterial effects of ethanol extracts from *Combretum quadrangulare* Kurz fruit collected in Kien Giang province, Vietnam. Research results showed that ethanol extract from *Combretum quadrangulare* Kurz fruit contained bioactive compounds such as alkaloids, flavonoids, steroids and triterpenoids, tannins, and phenolics. In the ethanol extract of *Combretum quadrangulare* Kurz fruit, the content of total phenolics and total flavonoids were recorded at 230.89 ± 2.36 mg GAE/g extract and 165.35 ± 5.80 mg QE/g extract, respectively. The survey results of in vitro antioxidant capacity through four methods of DPPH, ABTS⁺, RP and TAC indicated that the extracts of *Combretum quadrangulare* Kurz fruit showed its good activity with EC₅₀ values of 103.90 ± 0.40 µg/mL, 24.18 ± 0.06 µg/mL, 128.58 ± 0.20 µg/mL and 79.30 ± 1.40 µg/mL, respectively. The investigation results of in vivo antioxidant activity on fruit fly *Drosophila melanogaster* model showed that the fruit flies raised in the medium supplemented with 1 mg/mL extract of *Combretum quadrangulare* Kurz fruit had in vivo antioxidant effect under the conditions of oxidative stress induced by Paraquat and H₂O₂. In addition, the extract from *Combretum quadrangulare* Kurz fruit also had the ability to against the bacteria causing diseases in aquatic animals such as *Edwardsiella ictaluri*, *Aeromonas hydrophila* and *Streptococcus agalactiae*.

Index Terms antibacteria, antioxidant, phenolic, flavonoid, *Drosophila melanogaster*, *Combretum quadrangulare*

I. INTRODUCTION

Combretum quadrangulare Kurz (*C. quadrangulare*) is a plant species widely distributed throughout the Southern provinces of Vietnam. Most of them are cultivated, and some grow in the wild. In addition, *quadrangulare* is also found in Southeast Asian and African countries. It is commonly used in traditional medicine. Research on the chemical composition and biological activity of the compounds extracted from *quadrangulare* has reached remarkable world achievements. According to the statistics of Rebol [1], 97 compounds were extracted from *quadrangulare* in which 75 compounds belong to triterpenoids, and 19 compounds are in the group of flavonoids. These mentioned compounds belonged to the group of good antioxidant activity and antibacterial, anti-inflammatory, hepatoprotective and cytotoxicity abilities [2]. Seeds, leaves, and stem bark of *quadrangulare* were believed to have antipyretic and deworming effects [3].

Other studies have also found that the extracts from *quadrangulare* showed many good activities such as hepatoprotective activity [4], inhibitory ability against some bacterial species that caused diseases in aquatic animals [5]. A recent study by Dai et al. has concluded that the compounds isolated from the leaves of *quadrangulare* were effective against the bacterial strains of α-glucosidase and *Staphylococcus aureus* [6]. The study was carried out to find out the chemical composition, in vitro and in vivo antioxidant capacity, and the ability to resist pathogenic bacteria in aquatic animals of the ethanol extract of *quadrangulare* fruit. Thereby provide more scientific data on the biological activity of *C. quadrangulare* fruit, creating a premise for

the development of studies on screening the biological active compounds supporting and treating diseases in humans and aquatic animals species.

II. EXPERIMENTAL METHODOLOGY

A. Materials

C. quadrangulare fruit collected in Kien Giang province was identified by Dr. Nguyen Thi Kim Hue, Department of Biology, College of Natural Sciences, Can Tho University according to the Vietnamese plant taxonomy system.

Investigated bacterial strains: The research used pathogenic bacteria on aquatic species such as *Aeromonas dhakensis* (*A. dhakensis*), *Aeromonas hydrophila* (*A. hydrophila*), *Edwardsiella ictaluri* (*E. ictaluri*), and *Streptococcus agalactiae* (*S. agalactiae*). The mentioned strains were provided by the College of Aquaculture and Fisheries, Can Tho University, Vietnam.

Experimental fruit flies: Wild fruit fly *D. melanogaster* of Canton S (CS) strain was provided by Professor Kamei Kaeko, Kyoto Institute of Technology, Japan.

Chemicals Include ABTS (2,2-Azinobis 3-ethylbenzthiazolin-6-sulfonic acid), DPPH (2,2-Diphenyl-1-picrylhydrazyl), $K_2S_2O_8$, $K_3[Fe(CN)_6]$, $FeCl_3$, ammonium molybdate, Folin-Ciocalteu reagent, gallic acid, quercetin, paraquat, H_2O_2 , ethanol, methanol, and some other chemicals.

B. Extract Preparation

Removed the damaged parts of plant samples. After washing, the samples were dried, ground into small pieces, and soaked in ethanol 50 times for 24 hours each time. The extracts from the soaked were collected and filtered through filter paper. The extract was then subjected to evaporation to obtain the sample for analysis in further experiments.

C. Qualification of Natural Compounds

The extracts of *C. quadrangulare* fruit were qualitatively screened for their chemical components basing on their chemical properties as described by Jasuja et al. [7] with some adjustments. The data is presented in Table I.

TABLE I. QUALITATIVE SCREENING OF CHEMICAL COMPOSITION OF THE ETHANOL EXTRACT FROM *C. QUADRANGULARE* FRUIT

Chemical compound	Reagent	Identification
Alkaloids	Mayer	Orange brown to red precipitate
Flavonoids	Concentrated H_2SO_4	Dark yellow to orange, red precipitate
Steroids and Triterpenoids	Liebermann Burchard	Change to blue, green, orange, or red solution
Saponins	Distilled water and ethanol	Durable foaming
Tanins	Gelatin 1%	White cotton precipitate
Phenolics	$FeCl_3$ 10%	Blue-black or red orange precipitate

D. Quantification of Total Phenolic and Flavonoid Contents

Quantification of total phenolic content: Total phenolic content was determined according to the method described by Singleton et al. [8] with some adjustments. The reaction mixture of 250 μ L of extract, 250 μ L of distilled water, and 250 μ L of Folin-Ciocalteu reagent (1:4) was shaken well and incubated for 5 min at room temperature in the dark. Then, 250 μ L of Na_2CO_3 10% was added to the mixture and continued to incubate for 30 min at 40°C. The estimation of spectral absorbance of the reaction mixture was carried out at 765 nm. The total phenolic content was determined based on the gallic acid standard curve equation. Results were introduced in milligrams of gallic acid equivalent per gram of extract (mg GAE/g extract).

Quantification of total flavonoid content: The analysis of total flavonoid content was performed on the basis of the method described by Bağcı et al. [9] with some adjustments. The reaction mixture preparing with 200 μ L of extract, 200 μ L of distilled water, and 40 μ L of $NaNO_2$ 5%, was shaken and allowed to stand for 5 min. 40 μ L of $AlCl_3$ 10% was then added to the mixture and kept shaking well and staying still for 6 min. Then, the reaction mixture was added 400 μ L of 1M NaOH and 120 μ L of distilled water. The spectral absorbance of the reaction mixture was measured at 510 nm. The total flavonoid content was determined based on the quercetin standard curve equation. Results were displayed in milligrams of quercetin equivalent per gram of extract (mg QE/g extract).

E. Evaluation of in Vitro Antioxidant Activities

The investigation by DPPH method: The antioxidant activity of the extract was performed as the description in the method of Sharma and Bhat [10] with some adjustments. The reaction mixture of 100 μ L DPPH and 100 μ L extract was incubated in the dark for 60 min at room temperature. The measurement of the mixture absorbance spectra was implemented at 517 nm. The positive control used was gallic acid.

The investigation by ABTS method: The antioxidant activity of the extract was performed according to the method introduced by Nenadić et al. [11] with some adjustments. The reaction mixture consisting of 990 μ L $ABTS^{•+}$ and 10 μ L extract was incubated for 6 min at room temperature in the dark. Measured the absorbance spectra of the reaction mixture at 734 nm. The positive control used was gallic acid.

The investigation by Reducing Power (RP): The antioxidant activity of the extract was determined by the iron reduction capacity the method of Padmavathi et al. [12] with some adjustments. The reaction mixture containing 500 μ L of extract, 500 μ L of phosphate buffer (0.2 M, pH = 6.6), and 500 μ L of $K_3Fe(CN)_6$ 1% was incubated at 50°C for 20 min. Then, the mixture was added 500 μ L CCl_3COOH 10% and carried out the centrifugation at 3000 rpm for 10 min. Gently suck out 500 μ L of the

supernatant to add to 500 μ L of distilled water and 100 μ L of $FeCl_3$ 0.1%. Measured the absorbance spectra of the reaction mixture at 700 nm. The positive control applied was gallic acid.

The investigation by TAC method The antioxidant activity of the extract was performed by the method of Prieto et al. [13] with some adjustments. A mixture of 100 μ L extract and 900 μ L reagent solution was incubated at 95 $^{\circ}$ C for 90 min. The absorbance measurement was performed at 695 nm. The positive control was ascorbic acid.

F. In Vivo Antioxidant Activity

CS wild-type *D. melanogaster* played the role of research model in the study. The flies were reared at 25 $^{\circ}$ C under 12:12 light-dark cycle conditions. The in vivo antioxidant activity of the extract from *C. quadrangulare* fruit was tested in the fruit fly model as described by Etun et al. [14]. The experiment was carried out by selecting 48 h-closed male fruit flies to rear in three different food conditions, including medium supplemented with the extracts (1 mg/mL of feed), positive antioxidant medium (0.05 mg/mL gallic acid of feed), and standard medium. Fruit flies were replaced with new food every 2 days of the experiment. After 20 days, the fruit flies were kept in a fasted state for 5 hours, then transferred into test vials with 20 mM paraquat paper or O_2 10%. The experiment was reiterated 5 times for each treatment of 20 fruit flies. The number of survival flies was recorded after every 4 hours of observation during the experiment. The investigating indicators in the investigation included mean lifespan, 50% survival, and maximum lifespan. The maximum lifespan was calculated as the average survival time of 10% remaining in each test vial).

G. Antibacterial Assay

The antibacterial activity of the extracts was investigated by the agar plate diffusion method as described by Dast al. [15]. The antibacterial ability of the extract was determined based on the formation of an inhibition zone around the agar well. Inoculum with a concentration of 10^6 CFU/mL was spread evenly on the surface of Tryptecasein Soy Agar (TSA) plates with a volume of 100 μ L and allowed to dry for 15 min. Then, a small well of 6 mm in diameter was formed with sterile perforator, and 50 μ L of different extract concentrations prepared in DMSO 10% was added to the wells. The control treatment used DMSO 10%. Each extract concentration was repeated 3 times. After incubation for 24 h at 32 $^{\circ}$ C, the antibacterial ring diameter was determined as the circular inhibition zone forming around each well measured in mm.

Calculation of Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC): The minimum inhibitory concentration is identified as the lowest concentration that inhibits bacterial growth after 24 h of culture. The minimum inhibitory concentration was determined by the method of serial dilution of concentration in the tubes. Used a well culture plate,

each well contained 100 μ L of inoculation with a concentration of 10^6 CFU/mL and 100 μ L of the extracts at different concentrations. The wells were incubated aerobically at 32 $^{\circ}$ C for 24 hours. After incubation, the lowest concentration in the concentration range that inhibited bacterial growth was recognized as the minimum inhibitory concentration [16].

III. RESULTS AND DISCUSSION

A. Qualitative Analysis Results of Natural Compounds

The chemical constituents present in plant extracts are important for their biological activity. Therefore, determining the presence of chemical compounds is the first step of biological studies. The qualitative screening results of the chemical composition of the ethanol extract from *C. quadrangulare* fruit were presented in Table II.

TABLE II. CHEMICAL COMPOSITION PRESENTED IN THE ETHANOL EXTRACT OF *C. QUADRANGULARE* FRUIT

Chemical compound	Presence
Alkaloids	+
Flavonoids	+
Steroids and Triterpenoids	+
Saponins	-
Tanins	+
Phenolics	+

Note: (+) Present; (-) Absent

The results of Table II pointed out that the chemical components in *C. quadrangulare* fruit extract were determined to contain alkaloids, flavonoids, steroids and triterpenoids, tannins, and phenolics, based on the positive expression of the chemical reactions. However, the saponin group was not detected in the extract of *C. quadrangulare* fruit. It was possible that the presence of saponin compounds was not high enough to be identified in this study.

B. Quantification of Total Polyphenol and Flavonoid Contents

The total polyphenol content in the ethanol extract from *C. quadrangulare* fruit was determined based on the linear regression equation of gallic acid in the concentration range from 2 to 30 μ g/mL with $y = 0.082x + 0.0725$ ($R^2 = 0.9961$). The total flavonoid content in the extract was determined based on the linear regression equation of quercetin in the concentration range from 20 to 100 μ g/mL with $y = 0.0064x - 0.004$ ($R^2 = 0.9995$). The survey results showed that the total phenolic and flavonoid content of the ethanol extract of *C. quadrangulare* fruit were 230.89 ± 2.36 mg GAE/g extract and 165.35 ± 5.80 mg QE/g extract, respectively. Many studies have indicated that phenolic and flavonoid compounds have good biological activities. Research by Soobrattee et al. [17] demonstrated that phenolic compounds had redox properties; thus, they could act as antioxidants. Flavonoids found in plants had *in vitro* antioxidant activity and acted as *in vivo* antioxidants [18]. Besides, the antibacterial activity in phenolics and flavonoids was also determined [19]. All in all, it could

be reported that the extract of *C. quadrangulare* fruit contained natural compounds that have been introduced to have antioxidant and antibacterial activities.

C. In Vitro Antioxidant Activity

Plants are capable of producing secondary compounds including phenolics and flavonoids, which are the groups of compounds having antioxidant activity due to their

redox properties and chemical structure. The in vitro antioxidant effect of *C. quadrangulare* fruit extract was evaluated by determining the EC_{50} value of the extract in each method and comparing it with gallic acid. The smaller the EC_{50} value, the stronger the oxidation resistance. EC_{50} values of the extracts with standards are shown in Table III.

TABLE III. EC_{50} VALUES OF THE EXTRACTS AND THE STANDARD BY METHODS

Methods	Sample	Linear regression equation	EC_{50} ($\mu\text{g/mL}$)
DPPH	Gallic acid	$y = 12.10x + 0.3985$ ($R^2 = 0.9918$)	4.10 ± 0.10
	<i>C. quadrangulare</i>	$y = 0.448x + 3.4115$ ($R^2 = 0.9807$)	103.90 ± 0.40
ABTS ^{•+}	Gallic acid	$y = 90.16x + 4.8166$ ($R^2 = 0.9869$)	0.501 ± 0.002
	<i>C. quadrangulare</i>	$y = 1.821x + 5.9552$ ($R^2 = 0.9807$)	24.18 ± 0.06
RP	Gallic acid	$y = 0.050x - 0.0235$ ($R^2 = 0.9903$)	10.31 ± 0.10
	<i>C. quadrangulare</i>	$y = 0.003x + 0.0243$ ($R^2 = 0.9885$)	128.58 ± 0.20
TAC	Gallic acid	$y = 0.015x + 0.1206$ ($R^2 = 0.9863$)	25.34 ± 0.15
	<i>C. quadrangulare</i>	$y = 0.003x + 0.0243$ ($R^2 = 0.9885$)	79.30 ± 1.40

The results of Table III indicated that the in vitro antioxidant capacity of *C. quadrangulare* fruit extract in the 4 methods was lower than that of gallic acid. Specifically, the DPPH free radical neutralization effect of *C. quadrangulare* fruit extract had an EC_{50} value of $103.90 \mu\text{g/mL}$, higher than that of the standard of gallic acid ($EC_{50} = 4.10 \mu\text{g/mL}$). Therefore, the ability to scavenge 50% of DPPH free radicals of the extract was 25.34 times lower than that of gallic acid. Similarly, the ABTS^{•+} free radical scavenging efficiency of *C. quadrangulare* fruit extract had an EC_{50} value of $24.18 \mu\text{g/mL}$. In RP and TAC methods the EC_{50} value of the extract was $128.58 \mu\text{g/mL}$ and $79.30 \mu\text{g/mL}$, respectively. The values showed that the antioxidant activity of *C. quadrangulare* fruit extract was lower than that of the standard. Research by Nopsiri al. [20] demonstrated that the extracts from *C. quadrangulare* leaves with different solvents including methanol, hexane, and dichloromethane, had the ability to neutralize DPPH and ABTS^{•+} free radicals. Another plant species of the same genus, *Combretum tanaense* reported its antioxidant activity by Jaredet al. [21] through the DPPH survey method. The lowest value of EC_{50} means the highest antioxidant activity. Sample, which had EC_{50} or IC_{50} lower than $50 \mu\text{g/mL}$ was a very strong antioxidant; $50 - 100 \mu\text{g/mL}$ was a strong antioxidant and $100 - 150 \mu\text{g/mL}$ was a medium antioxidant while a weak antioxidant with IC_{50} or $EC_{50} > 150 \mu\text{g/mL}$ [22]. Accordingly, EC_{50} values in this study ranged from 24.18 to $128.58 \mu\text{g/mL}$, showing that *C. quadrangulare* fruit extract expressed good antioxidant activity under in vitro conditions.

D. The Effect of in Vivo Antioxidant Activity

The antioxidant capacity of *C. quadrangulare* fruit extract was tested in vivo on the fruit fly model through its effectiveness against oxygen stress under H_2O_2 -induced conditions. The in vivo antioxidative effect of fruit flies reared in the medium supplemented with gallic acid of 0.05 mg/mL and *C. quadrangulare* fruit extract of 1 mg/mL under oxidative stress induced by 20 mM PQ and $H_2O_2 10\%$ was introduced (Fig. 1 and Fig.

2. The three values expressed the results, including mean lifespan, 50% survival, and maximum lifespan.




Figure 1. In vivo antioxidant effect of *C. quadrangulare* fruit extract under PQ condition.




Figure 2. In vivo antioxidant effect of *C. quadrangulare* fruit extract under H₂O₂ condition.

The results concluded that under the conditions of oxidative stress caused by 20 mM PQ and $H_2O_2 10\%$, the

survival ability of fruit flies reared in the medium supplemented with gallic acid and extract of *C. quadrangulare* fruit was longer than that of the fruit flies reared in standard medium. Specifically, the mean lifespan, 50% survival, and maximum lifespan of fruit flies raised in feed supplemented with extracts of *C. quadrangulare* fruit under PG-induced oxidative stress were 22.53 h, 19.00 h, and 45.00 h, respectively, which meant higher than that of the control of 1.30 times, 1.24 times and 2.02 times, respectively (Fig. 1). Similar results were also observed in H₂O₂-induced oxidative stress, where the mean lifespan, 50% survival, and maximum lifespan were 39.67 h, 38.00 h, and 53.33 h, respectively. The data were higher than the control at 1.33, 1.16, and 1.34 times, respectively (Fig. 2). The research results once again contributed to confirming the antioxidant potential of the extract of *C. quadrangulare* fruit. This result was consistent with the qualitative, quantitative, and in vitro antioxidant survey results. It is known that fruit flies are the model used in many studies *in vivo* antioxidant activity [23, [24]. Therefore, the experiment results are reliable.

E. Antibacterial Activity

The in vitro antibacterial activity of the ethanol extract of *C. quadrangulare* fruit is presented in Table IV. The extract had antibacterial activity against Gram positive and Gram negative bacteria, with MIC values ranging from 320 to 1280 mg/mL. The results in Table V showed that *C. quadrangulare* extract was resistant to three strains of pathogenic bacteria in aquatic animals, including *A. hydrophila*, *E. ictaluri* and *S. agalactiae*. However, *C. quadrangulare* extract was not resistant to *A. dhakensis*.

TABLE IV. ANTIBACTERIAL ACTIVITY OF ETHANOLIC EXTRACT OF *C. QUADRANGULARE* FRUIT

Bacteria strains	Inhibition zone (mm)						MIC (mg/mL)
	40	80	160	320	640	1280	
<i>A. dhakensis</i>	-	-	-	-	-	-	-
<i>A. hydrophila</i>	-	2.5	3.8	4.4	5.8	7.0	640 < MIC < 1280
<i>E. ictaluri</i>	-	3.0	4.6	6.0	7.2	9.7	320 < MIC < 640
<i>S. agalactiae</i>	-	3.1	5.2	6.3	7.6	10.3	320 < MIC < 640

Note: (-) not resistant

Research results of Hant al. [5] showed that the extracts of *C. quadrangulare* seeds and leaves were resistant to pathogenic bacteria on aquatic animals such as *Aeromonas hydrophila*, *Edwardsiella ictaluri* and *Vibrio parahaemolyticus*. In addition, the methanol extract of *C. quadrangulare* seeds was resistant to *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Acinetobacter baumannii* and *Escherichia coli* strains with the MIC values of 1212 g/mL, 2425 g/mL, 2425 g/mL, and 9700 g/mL, respectively [25]. Phenolic and flavonoid compounds have been previously reported to have antifungal and antibacterial activities [26, 27]. Other research has also shown that flavonoids and phenolics are common secondary compounds synthesized in plants; besides that, these compounds have been studied and shown to have antibacterial effects [28]. In this study, the amount of phenolic and flavonoid identified in *C. quadrangulare* fruit extract was 230.89 ± 2.36 mg GAE/g extract and 165.35 ± 5.80 mg QE/g extract, respectively. Therefore, the antibacterial activity shown in this experiment is possibly due to the presence of phenolic and flavonoid groups in the extract.

IV. CONCLUSION

Research results showed that ethanol extract of *C. quadrangulare* fruit contained biologically active compounds such as alkaloids, flavonoids, steroids and triterpenoids, tannins, and phenolics. The investigated extract also expressed the *in vitro* and *in vivo* antioxidant activity. In addition, the activities against pathogenic bacteria strain on aquatic animals such as *A. hydrophila*, *E. ictaluri* and *S. agalactiae* were also recognized in the extract. The research results provided evidence that *C. quadrangulare* fruit extract was a potential herb

containing many natural compounds with good biological activity. The study provided valuable scientific information for further studies in finding solutions to apply *Combretum quadrangulare* Kurz in pharmacology and disease prevention for aquatic animals.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest with the contents of this article.

AUTHOR CONTRIBUTIONS

Tran Thanh Men and Nguyen Trong Tuan were responsible for the study design; Tran Thanh Mer was responsible for study design and manuscript writing; Huynh Kim Yen, Huynh Thi Cam Lan and Nguyen Hoang Son conducted the practical experimental work; Tan Khang made a statistical analysis of the experiment. All authors had approved the final version.

ACKNOWLEDGMENT

We are grateful to Professor Kaeko Kamei, Department of Biomolecular Engineering, Kyoto Institute of Technology, Japan, for providing the *Drosophila melanogaster* Canton-S.

REFERENCES

[1] R. Roy, R. K. Singh, S. K. Jash, A. Sarkar and D. Gorai, "Combretum quadrangulare (Combretaceae): Phytochemical constituents and biological activity," *Indo Am. J. Pharm. Resol.* 4, no. 8, pp. 3413-3427, 2014.
 [2] V. Sharma, S. Sharma and R. P. Pracheta, "Withania somnifera A rejuvenating ayurvedic medicinal herb for the treatment of

- various human ailments Int. J. Pharmtech Resvol. 3, no. 1, pp. 187-192, 2011.
- [3] A. H. Banskota, Y. Tezuka, Q. Tram and S. Kadota, "Chemical constituents and biological activities of Vietnamese medicinal plants" Curr. Top. Med. Chemvol. 3, no. 2, pp. 222-248, 2003.
- [4] I. K. Adnyana, Y. Tezuka, A. H. Banskota, K. Q. Tram and S. Kadota, "Three new triterpenes from the seeds of Combretum quadrangulare and their hepatoprotective activity" J. Nat. Prod, vol. 64, no. 3, pp. 366-363, 2001.
- [5] T. T. T. Hang, C. T. Duc, L. T. T. Vy and N. C. Trang, "Study on the antibacterial activities of extracts from sakae nao (Combretum quadrangulare) on diseased aquatic animals and bacteria under in vitro conditions" Can Tho Univ. J. Sci.vol. 13, pp. 106112, 2018.
- [6] T. B. N. Dao, T. M. T. Nguyen, V. Q. Nguyen, et al, "Flavones from Combretum quadrangulare growing in Vietnam and their-glucosidase inhibitory activity" Molecules vol. 2021, no. 26, .p 2531, 2021.
- [7] N. D. Jasuja, S. K. Sharma, R. Saxena, et al, "Antibacterial, antioxidant and phytochemical investigation of Thuja orientalis leaves" J. Med. Plant Resvol. 7, no. 25, pp. 1886-1893, 2013.
- [8] V. L. Singleton, R. Orthofer and R. M. Lamuela-Raventós, "Analysis of total phenol and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent" Methods Enzymol, vol. 299, pp. 152-178, 1999.
- [9] G. C. Bag, P. G. Dey and T. H. Bhaigyaabati, "Assessment of total flavonoid content and antioxidant activity of methanolic rhizome extract of three Hedychium species of Manipur valley" Int. J. Pharm. Sci, vol. 30, no. 1, pp. 154-159, 2015.
- [10] O. P. Sharma and T. K. Bhat, "DPPH antioxidant assay revisited," Food Chem.vol. 113, no. 4, pp. 1202-1205, 2009.
- [11] N. Nenadis, L. F. Wang, M. Tsimidou et al, "Estimation of scavenging activity of phenolic compounds using the ABTS assay" J. Agric. Food. Chem.vol. 52, no. 15, pp. 4666-4674, 2004.
- [12] R. Padma, N. G. Parvathy, V. Renjith et al, "Quantitative estimation of tannins, phenols and antioxidant activity methanolic extract of Imperata cylindrica" Int. J. Pharm. Sci. Res, vol. 4, no. 1, pp. 73-77, 2013.
- [13] P. Prieto, M. Pineda and M. Aguilar, "Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: Specific application to the determination of vitamin E Anal. Biochem, vol. 269, no. 2, pp. 337-341, 1999.
- [14] M. A. Etuh, J. C. Aguiyi S. O. Ochala, et al, "The in vivo antioxidant protective activity of Mangifera indica cold aqueous leaf extract in Drosophila melanogaster" J. Adv. Biol. Biotechnol. vol. 22, no. 2, pp1-7, 2019.
- [15] T. Das, S. Seha and M. Manefield, "The roles of extracellular DNA in the structural integrity of extracellular polymeric substance and bacterial biofilm development" Environ. Microbiol. Rep, vol. 5, no. 6, pp. 778-786, 2013.
- [16] J. G. Usman, O. A. Sodipo and U. K. Sandabe, "In vitro antimicrobial activity of Cucumis metuliferu E. Mey. Ex. Naudin fruit extracts against Salmonella gallinarum" Int. J. Phytomedicinevol, 6, no. 2, pp. 262-274, 2014.
- [17] M. A. Soobrattee, V. S. Neergheen, A. Luximonamma, et al, "Phenolics as potential antioxidant therapeutic agents: Mechanism and actions" Mutat. Res. Fund. Mol. Mech. Mutagenol. 579, no. 1-2, pp. 209-213, 2005.
- [18] S. Geetha, M. S. Ram and S. S. Mongia et al, "Evaluation of antioxidant activity of leaf extract of Seabuckthorn Hippophae rhamnoides L.) on chromium (VI) induced oxidative stress in albino rats" J. Ethnopharmacol.vol. 87, no. 23, pp. 247-251, 2003.
- [19] R. Baharfar, R. Azimi and M. Mohseni, "Antioxidant and antibacterial activity of flavonoid, polyphenol and anthocyanin rich extracts from Thymus kotschyano-boiss & hohen aerial parts" J. Food Sci. Technolvol. 52, no. 10, pp. 6777-6783, 2015.
- [20] W. Nopsiri, S. Chansakaow, S. Putiyanan, et al, "Chemical constituents and antibacterial activity of volatile oils of Combretum latifolium Bl. and C. quadrangulare Kurz Leaves" Chiang Mai Univ. J. Nat. Sci.vol. 14, pp. 245-256, 2015.
- [21] M. O. Jared, A. W. Bibiane, A. M. Gervasoni et al., "The antibacterial antioxidant and phytochemical composition of Combretum tanaense (d. Clark) root extracts" Eur. J. Med. Chem. vol. 23, no. 4, pp. -8, 2018.
- [22] S. Kusmardiyani, G. Novita and I. Fidrianny, "Antioxidant activities from various extracts of different parts of kelakai (Stenochlaena palustis) grown in central Kalimantan – Indonesia" Asian J. Pharm. Clin. Resvol. 9, pp. 2152-219, 2016.
- [23] E. Bier, "Drosophila the golden bug, emerges as a tool for human genetics." Nat. Rev. Genetvol. 6, no. 1, pp. 23, 2005.
- [24] V. Padalko, V. Dzyuba, O. Kozlova et al., "Zingiber officinale extends Drosophila melanogaster life span in xenobiotic induced oxidative stress conditions" Front. Biol., vol. 13, no. 2, pp. 130-136, 2018.
- [25] K. Natachit, D. Santiamorn, and B. Khantawa, "Antibacterial activity of the seeds of Combretum quadrangulare Kurz (Combretaceae) Chiang Mai J. Sci.vol. 5, no. 3, pp. 333-339, 2006.
- [26] W. F. Zheng R. X. Tan and L. Yang et al, "Two flavones from Artemisia giraldii and their antimicrobial activity" Planta Med, vol. 62, no. 2, pp. 160-162, 1996.
- [27] A. J. Afolayan and J. J. Meyer, "The antimicrobial activity of 3,5,7-trihydroxyflavone isolated from the shoots of Melichrysum aureonitens" J. Ethnopharmacol.vol. 57, no.3, pp. 177-181, 1997.
- [28] S. A. Baba and S. A. Malik, "Determination of total phenolic and flavonoid content, antimicrobial and antioxidant activity of a root extract of Arisaema jacquemontii Blume," J. Taibah Univ. Sci. vol. 9, no. 4, pp. 444-454, 2015.



Tran Thanh Men is currently a senior lecturer of the Biology Department under the College of Natural Sciences in Can Tho University, Viet Nam. He obtained his Ph.D. in Material and Life science at Kyoto Institute of Technology, Kyoto city, Japan in 2016. Tran Thanh Men's early studies were carried out from 2003 in the biomolecular field. In 2007 he participated in research on malaria and medicinal plants with antimalarial activity. In 2012, he took part in research on

the fruit fly *Drosophila melanogaster* and the genes involved in the metabolic processes of the fruit flies. Since 2017, he has spent time on various studies on plant species with weed resistance and insecticidal properties and those with anti-aging activity.